



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Reduced Cancer Mortality at High Altitude: The Role of Glucose, Lipids, Iron and Physical Activity

Thiersch, M ; Swenson, E R ; Haider, T ; Gassmann, M

Abstract: Residency at high altitude (HA) demands adaptation to challenging environmental conditions with hypobaric hypoxia being the most important one. Epidemiological and experimental data suggest that chronic exposure to HA reduces cancer mortality and lowers prevalence of metabolic disorders like diabetes and obesity implying that adaption to HA modifies a broad spectrum of physiological, metabolic and cellular programs with a generally beneficial outcome for humans. However, the complexity of multiple, potentially tumor-suppressive pathways at HA impedes the understanding of mechanisms leading to reduced cancer mortality. Many adaptive processes at HA are tightly interconnected and thus it cannot be ruled out that the entirety or at least some of the HA-related alterations act in concert to reduce cancer mortality. In this review we discuss tumor formation as a concept of competition between healthy and cancer cells with improved fitness - and therefore higher competitiveness - of healthy cells at high altitude. We discuss HA-related changes in glucose, lipid and iron metabolism that may have an impact on tumorigenesis. Additionally, we discuss two parameters with a strong impact on tumorigenesis, namely drug metabolism and physical activity, to underpin their potential contribution to HA-dependent reduced cancer mortality. Future studies are needed to unravel why cancer mortality is reduced at HA and how this knowledge might be used to prevent and to treat cancer patients.

DOI: <https://doi.org/10.1016/j.yexcr.2017.03.048>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-136692>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Thiersch, M; Swenson, E R; Haider, T; Gassmann, M (2017). Reduced Cancer Mortality at High Altitude: The Role of Glucose, Lipids, Iron and Physical Activity. *Experimental Cell Research*, 356(2):209-216.

DOI: <https://doi.org/10.1016/j.yexcr.2017.03.048>

Reduced Cancer Mortality at High Altitude: The Role of Glucose, Lipids, Iron and Physical Activity

M.Thiersch^{1,*}, E. R. Swenson^{2,3}, T. Haider¹ and M. Gassmann^{1,4}

Affiliations: ¹Institute of Veterinary Physiology, Vetsuisse Faculty, and Zurich Center for Integrative Human Physiology (ZIHP), University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland. ²Dept of Medicine, Division of Pulmonary and Critical Care Medicine, University of Washington, Seattle, WA, USA. ³Medical Service, Veterans Affairs Puget Sound Health Care System, Seattle, WA, USA. ⁴Universidad Peruana Cayetano Heredia (UPCH), Lima, Peru.

*Corresponding author: Markus Thiersch, ¹Institute of Veterinary Physiology, Vetsuisse Faculty, and Zurich Center for Integrative Human Physiology (ZIHP), University of Zurich, Winterthurerstrasse 260, 8057 Zurich, Switzerland. E-mail:markus.thiersch@uzh.ch

1	Introduction	1
2	Epidemiology.....	2
2.1	Human Studies.....	2
2.2	Animal Studies	2
3	Cancer and Adaptation.....	3
3.1	Metabolic Alterations.....	4
3.1.1	Glucose and Glutamine Metabolism.....	4
3.1.2	Lipid Metabolism.....	5
3.1.3	Iron Metabolism and Anemia of Cancer	6
3.2	Pharmacokinetics and Pharmacodynamics at HA	8
3.3	Physical Activity	9
4	Conclusion.....	11
5	Acknowledgements	12
6	References	13

Abstract

Residency at high altitude (HA) demands adaptation to challenging environmental conditions with hypobaric hypoxia being the most important one. Epidemiological and experimental data suggest that chronic exposure to HA reduces cancer mortality and lowers prevalence of metabolic disorders like diabetes and obesity implying that adaption to HA modifies a broad spectrum of physiological, metabolic and cellular programs with a generally beneficial outcome for humans. However, the complexity of multiple, potentially tumor-suppressive pathways at HA impedes the understanding of mechanisms leading to reduced cancer mortality. Many adaptive processes at HA are tightly interconnected and thus it cannot be ruled out that the entirety or at least some of the HA-related alterations act in concert to reduce cancer mortality. In this review we discuss tumor formation as a concept of competition between healthy and cancer cells with improved fitness – and therefore higher competitiveness – of healthy cells at high altitude. We discuss HA-related changes in glucose, lipid and iron metabolism that may have an impact on tumorigenesis. Additionally, we discuss two parameters with a strong impact on tumorigenesis, namely drug metabolism and physical activity, to underpin their potential contribution to HA-dependent reduced cancer mortality. Future studies are needed to unravel why cancer mortality is reduced at HA and how this knowledge might be used to prevent and to treat cancer patients.

1 Introduction

Humans living at moderate (above 1500 m) and high (above 2500 m) altitude (HA) are chronically exposed to hypobaric hypoxia and need to adapt to this environmental challenge. The altitude-dependent drop in barometric pressure and subsequent decrease of the ambient partial pressure causes hypoxemia and tissue hypoxia. When sojourners acclimatize, their bodies respond first with short-term adaptations (hours to days; e.g. increased basal ventilation [1], plasma volume reduction [2, 3]) and later with long-term adaptations (weeks to months; e.g. increased erythropoiesis [4]) that are mainly driven by tissue hypoxia. The magnitude of response to HA is the key factor for physiological adaption or pathological mal-adaption, the latter resulting in hypoxia-related altitude illnesses [5-7]. The cellular response to hypoxia is driven by hypoxia inducible transcription factors (HIF) that are rapidly activated when intracellular oxygen levels decline [8-10]. Iron-dependent prolyl-hydroxylases (PHDs) regulate the stability of HIF- α subunits resulting in either protein degradation when oxygen is sufficiently available or stabilization when oxygen supply is low [11]. The detailed regulation of HIFs by oxygen availability and their down stream targets have been extensively reviewed [11-13]. During oxygen deficiency, hematopoiesis is driven by the renal hormone erythropoietin (Epo) that is oxygen-dependently regulated by HIF-2 [14] and promotes red blood cell production.

Epidemiological studies provide evidence that humans populating high altitude environments show reduced cancer mortality. Several possible explanations have been discussed in the past including elevated vitamin D synthesis due to higher background radiation [15]. However, the most adaptive processes at HA occur in response to reduced oxygen availability and this might be a much greater factor in cancer mortality. At the first glance, the observation of oxygen-dependent reduced cancer mortality at HA is counterintuitive because tumor hypoxia *per se* is rather a supportive factor for tumor growth and the development of aggressive phenotypes [16, 17]. Both hypoxia-inducible factors 1 and 2 (HIF-1/2) have been reported to promote tumor growth and metastasis [18-20]. In contrast, some studies show that the loss of HIF-1 α expression promotes kidney cancer [21] suggesting that at least in some tissues

stabilized HIF is tumor suppressive. Recent studies analyzed evolutionary selected variants *EGLN1* (encoding for PHD2) and *EPAS1* (encoding for HIF-2 α) genes in Tibetans and reported association of these variants with increased lung [22] and gastric [23] cancer risk. These HA-adapted variants of *EGLN1* with D4E and C127S polymorphisms increased PHD2 activity and promote HIF- α degradation [24, 25] suggesting that HIF stabilization at HA might contribute to reduced incidence/mortality of at least some types of cancer. In this review we briefly summarize the results of epidemiological and animal studies and discuss mainly oxygen-based mechanisms that may account for reduced cancer mortality. It should be mentioned that “mortality” is a clinical relevant outcome but does not further explain at which steps during the course of cancer disease (incidence, progression, dissemination, response to therapy) exposure to HA has affected and altered.

2 Epidemiology

2.1 Human Studies

In 1974/75 an inverse relationship of HA and leukemia cancer mortality in humans was reported for the first time [26-28]. These studies were followed by additional studies within the last decades that considered potential cofounders (such as environment and socio-economic status) and confirmed these findings [29-35]. Although overall cancer mortality negatively correlates with HA [31], site-specific cancer seem to respond differently to HA. While lymphoma, breast [36], lung, cancers of tongue and mouth or larynx [29, 35] display reduced mortality at HA, liver and cervix mortality showed no difference between low and high altitude. As expected, mortality from melanoma was increased due to the higher background radiation [37]. Very recently, it has been shown that highlanders display a reduced incidence of lung and breast cancer [38] supporting earlier observations that HA is indeed involved in prevention of tumorigenesis.

2.2 Animal Studies

Reduced cancer mortality at HA is not unique to humans but has been observed in mice exposed to very high altitude (4540 m) for a prolonged period of time after exposure to sub-

lethal levels of x-rays resulting in reduced incidence of tumor formation [39]. Additionally, the incidence of spontaneous leukemia is reduced at HA [40] but spontaneously forming pulmonary tumors display a higher incidence at HA [41]. Back in the 1980s, studies in several rat and mouse tumor models demonstrated reduced primary tumor growth and metastatic spread as well as improved response to therapeutic treatment at HA [42-44]. A more recent study showed reduced tumor incidence not only in two spontaneously developing tumor mouse models, namely $p53^{-/-}$ and $APC^{Min/+}$, but also in a chemically induced skin cancer model in mice exposed to 10% normobaric oxygen [45] corresponding to an extreme altitude of approximately 5500 – 6000 meter above sea level [12]. With their experimental setup, the latter authors prove that reduced oxygen levels protect from oncogenic events and tumorigenesis matching the epidemiological findings of Simeonov and Himmelstein [38].

3 Cancer and Adaptation

Theoretically, cancer cells can be considered to be evolutionary selected cell lineages that have escaped control of replication and cell death and meet the traditional characteristics of minimal Darwinian populations, namely variation, selection and inheritance [46]. As a consequence, the evolutionary process of tumor formation, progression and metastasis depends on the nature of the niche that is populated by cancer cells and their ability to compete for resources with healthy cells within the same niche. Natural selection demands genotypes that differ in fitness to result in improved reproduction success. Although oncogenic events are commonly assumed to increase cellular fitness, DeGregori, (2011) suggests a model of adaptive oncogenesis, in which genetic variations rarely result in advantageous traits in healthy (and young) populations of (stem) cells, since they always display a high degree of fitness to maintain tissue integrity and this successfully competes against somatic cancer cell evolution [47]. Cell age and/or damaging insults reduce cellular fitness and change the adaptive landscape due to accumulation of mutations and alterations in the microenvironment, both promoting selection of adaptive oncogenesis events [47]. An example in relation to cancer at HA is melanoma that is increased due to increased radiation exposure [37] resulting in increased damage of epidermal cells and consequently the reduction in cell fitness. In the light of this hypothesis, we discuss below reduced cancer

mortality and incidence at HA as a competitive mechanism between cancer and healthy cells. We suggest that HA increases fitness of healthy cells and/or modifies the tumor microenvironment resulting in adverse conditions for tumor cell development and expansion. Other diseases (e.g. stroke and cardiovascular diseases) display also a link between high altitude and reduced mortality [48] supporting our assumption. As a matter of fact, the metabolic syndrome, diabetes type 2 and obesity are cancer risk factors [49, 50] and have a lower prevalence in humans populating elevated areas [51-53]. In the next sections, we consider cell biological and metabolic alterations at HA that might pose a bottleneck for emerging tumors in humans living at higher elevations.

3.1 Metabolic Alterations

3.1.1 Glucose and Glutamine Metabolism

The competition for nutrition in a resource-restricted environment may result in different metabolic pathways being susceptible to alterations that might occur at HA. The most prominent metabolic reprogramming in tumor cells is the “Warburg effect” resulting in aerobic metabolism of glucose and increased lactate production – even when sufficient oxygen is available [54]. A recent review summarizing tumor metabolism and discussing metabolites that limit tumor progression as potential therapeutic targets suggested that neither ATP nor NADPH may limit tumor proliferation [54]. However, restoration of tricarboxylic acid (TCA) cycle metabolites and the synthesis of nucleotides may be a limiting factor in tumor progression due to the high demand for DNA synthesis in proliferating cells. Glucose and/or glutamine (depending on tumor type and microenvironment) refuel the TCA cycle and provide substrates for the nucleotide metabolism [55, 56]. Consequently, both, glucose and glutamine may limit tumor progression when insufficiently supplied to cancer cells (e.g. when healthy cells increase their demand, or when less substrate is provided from blood). As expected, hyperglycemia correlates with an increased risk of tumorigenesis [57] and because highlanders have reduced plasma glucose and glutamine concentrations (at least in acute and prolonged exposure to HA [58, 59]) this restriction possibly contributes to reduced cancer mortality [58, 60]. Blood glucose levels are mainly controlled by the liver and partially by kidneys [61, 62] and release of hepatic glucose is stimulated by glucagon, fatty acids and

catecholamines but suppressed by insulin [61]. Accordingly, as Andean highlanders display higher glucagonemia and lower glycemia, their exposure to HA might reduce hepatic glucagon sensitivity resulting in reduced glucose release from the liver. In cancer cells, HIF-1 increases the expression levels of glucose transporter 1 and 3 (GLUT1/3) to facilitate glucose uptake [63]. However, adipocytes respond similarly to hypoxia with increased GLUT1/3 expression levels [64]. Furthermore, acclimatization to HA increases the uptake and metabolism of blood glucose by human skeletal and heart muscle [61], facilitated by a hypoxia-driven increase in HIF-1 as well as GLUT4 expression levels in human muscle cells [61, 65].

In conclusion, cancer cells are competing with healthy cells and tissues (e.g. muscles) for nutrition including glucose and glutamine and exposure to HA possibly limits the availability (reduced plasma levels) and increases glucose uptake and metabolism in healthy tissue (competition) for these metabolites. However, metabolic pathways are intricately interconnected and many cancer cells utilize a broad spectrum of metabolites including lipids, lactate and further amino acids [54] potentially compensating reduced availability of even central substrates like glucose.

3.1.2 Lipid Metabolism

Lipid metabolism is a complex process that is regulated by multiple pathways to control uptake, transport, metabolism and synthesis. In turn, lipids also differentially regulate cellular signaling transduction and pathways [66]. The cell membrane, mainly consisting of lipids, contains lipid rafts that are spatial membrane assemblies of cholesterol and derivatives providing a platform for signaling transduction processes involved in apoptosis [67] as well as cell survival and proliferation [68, 69]. Small variations in the tightly regulated metabolism of lipids result in disturbed energy homeostasis ultimately leading to fat accumulation, increased body mass and obesity, which are risk factors for breast [70], colon [71], kidney [72], prostate [73], gallbladder [74] and pancreas cancer [75]. Fast growing tumors display dysregulated lipid homeostasis with reduced 3-hydroxy-3-methyl-glutaryl CoA (HMG-CoA) reductase and low-density lipoprotein (LDL) receptor activity as well as cholesterol enrichment in tumor cells [76]. Cancer cells require energy as well as metabolic precursors for cell proliferation associated with rapid lipid synthesis for assembling membranes of new cells. Lipids are

provided from external sources as well as intracellularly by fatty acid synthesis [76] and regulate, besides membrane assembly, the cancer related pathways PI3K/AKT/mTOR, RAS and Wnt [76-78]. Further, free fatty acids in human plasma are also involved in insulin resistance [79] resulting in high insulin and insulin-like growth factors (IGF) that directly promote tumor progression [73, 76]. The lipid profile of tumors in patients at HA has not been published, but hypoxia has been shown to facilitate exogenous lipid uptake of cancer cells [80, 81]. Such a lipid supply might be suppressed in highlanders who have been reported to exhibit lower fatty acid plasma levels [61] and potentially decrease fatty acid metabolism due to SIRT4 induction [82]. Elevated lipid levels are commonly accepted to be severe risk factors for cancer [83, 84]. Highlanders living at moderate altitude (1500 m) have reduced levels of total cholesterol and triglycerides but an increase of high density lipoprotein (HDL) [85, 86] that correlates with reduced prevalence of both cardiovascular diseases [87, 88] and cancer incidence including breast [89], prostate [90, 91] and potentially colon cancer, as shown in mice [92]. Collectively, the restricted lipid metabolism at HA might be shifted against the development and progression of tumors.

3.1.3 Iron Metabolism and Anemia of Cancer

Iron is an essential nutrient involved in the formation of red blood cells, in many oxidation-reduction processes within the organism as well as in energy metabolism, mitochondrial respiration, and DNA synthesis. Iron metabolism is controlled on both the systemic and cellular levels. The liver hormone hepcidin is a key regulator for iron homeostasis inhibiting intestinal iron absorption as well as macrophage release by degrading the iron-transport protein ferroportin (Fpn1) [93-95]. Hypoxia is tightly associated with iron metabolism. HIFs control directly or indirectly the expression of the divalent metal transporter 1 (DMT-1) and Fpn1 among others [96, 97] and iron is an essential cofactor for PHDs that catalyze the hydroxylation of HIF- α subunits to target them for degradation [11-13]. Under hypoxic conditions hepcidin mRNA transcription is indirectly inhibited by Epo via erythroferrone [96, 98] to guarantee sufficient iron supply for red blood cell formation. Cancer cells have a higher iron demand and thus elevated iron uptake. Correspondingly, high iron levels have been determined as a risk factor for cancer [99-101] with an impact on tumor-associated macrophages and the tumor microenvironment [102]. Cancer cells display disturbed iron

homeostasis with increased expression of transferrin receptor 1 (uptake) [103], decreased expression of ferritin (storage) [104, 105] and inhibition of ferroportin (release) [106]. Exposure to HA demands increased hematopoiesis to cope with reduced oxygen uptake and thus is associated with elevated iron utilization for red blood cell formation [96]. At very high altitudes female adults develop reduced body iron levels [107] potentially decreasing the risk for tumor incidence. Acute or short-term exposure to HA decreases plasma iron levels, ferritin and transferrin saturation depending on the dietary iron intake [108, 109]. The critical role of iron in malignancy has been demonstrated by use of iron chelators that sequester iron or by applying antibodies against the transferrin receptor 1 [110-112]. Both approaches inhibit cancer progression. Iron deficiency is the most prevalent nutritional disorder [113] resulting in anemia. This includes anemia of cancer [114] that frequently occurs in patients secondary to their cancer [114] or is induced during chemo and/or radiotherapy [115, 116]. Anemia of cancer reduces quality of life and is a poor prognostic factor in cancer patients [117, 118]. Consequently, it demands treatment that is currently achieved by iron supplementation, administration of erythropoiesis-stimulating agents (ESA) and blood transfusion [114]. Hepcidin expression (the aforementioned key regulator of iron homeostasis) is frequently induced in tumor patients by cancer-induced inflammatory signals [119] and prevents iron uptake resulting in iron deficiency and finally anemia. Interestingly, hepcidin levels – at least during acute and prolonged exposure to HA and isobaric hypoxia – decline to meet the increased iron demand [120-123]. Epidemiological data on hepcidin expression levels and anemia of cancer in highlanders are missing and, consequently, thorough reports on frequency, severity and treatment success are required to determine its role in cancer mortality at HA. It has been reported, however, that the frequency of end-stage renal disease dependent anemia (which can occur in cancer patients as a result of renal tumor invasion [114]) is reduced [124, 125]. Additionally, the response to Epo or erythropoiesis-stimulating agents (ESA) is improved at HA [126, 127]. Besides iron deficiency, impaired Epo production (due to cancer- or therapy-induced kidney damage) is the major cause for anemia of cancer [114]. With increasing altitude the number of end-stage renal disease patients receiving ESA as well as the required dosages surprisingly are lower while hemoglobin levels as well as patient survival are higher [126, 127]. The reduced requirement of ESA treatment in anemic

patients at HA may also have implications for cancer mortality. Several clinical [128-135] and preclinical [136-144] studies suggest that ESA treatment in cancer patients actually reduces patient survival by promoting tumor growth via binding to Epo receptors on many cancer cell types. Hence, the FDA has lowered the minimally accepted hemoglobin levels in anemic cancer patients (lowlanders) to <12 g/dl [145, 146]. A potentially lower demand of ESA to treat anemia of cancer at HA might thus contribute to reduced cancer mortality at HA.

3.2 Pharmacokinetics and Pharmacodynamics at HA

A frequently ignored subject is the alteration of drug metabolism at HA. In addition to surgery and irradiation, chemotherapy is the most frequent choice of treatment in cancer. The pharmacokinetic and pharmacodynamic responses to drugs have been poorly studied in highlanders [147]. With respect to the approximately 400 million humans living at elevated areas (above 1500 m) [48, 148], this population is considerably underrepresented, especially because there is clear evidence that HA exposure (like all other physiological stress conditions) has impact on drug absorption, distribution, metabolism and detoxification and clearance [147]. Different types of chemotherapy used to treat cancer include alkylating agents, antimetabolites, anti-microtubule agents, topoisomerase inhibitors, antibiotics, and antibodies and all may have a potentially altered efficacy at HA. Anthracyclines (e.g. doxorubicin) or alkylating agents (cisplatin) generate high levels of reactive oxygen species (ROS) that have an impact on cancer cells [149]. Since ROS formation is a tightly balanced process, alterations during cancer therapy may be either tumor promoting (e.g. genetic instability) or inhibiting [150]. Exposure to HA alters the mitochondrial respiratory chain by increasing the COX4-2 expression in a HIF-1 mediated fashion. COX4 is a subunit of cytochrome oxidase and the alternative COX4-2 variant replaces COX4-1 during hypoxia, and so increasing the efficiency of electron transfer in complex IV under low oxygen concentrations [151]. This might prevent excessive ROS formation in cancer cells, but also, as in healthy muscle cells [152, 153], potentially protects organs and tissues from cytotoxic therapies. Furthermore, reduced expression of cytochrome P450 affects the metabolic clearance of drugs [154] and altered blood flow to organs might cause changes in drug distribution in comparison to lowlanders. It is also not clear if HA changes tumor perfusion

and thereby the tumor accessibility of the drugs. The response to tumor therapies at high altitude with and without therapy-associated anemia would be a very interesting area to focus future research efforts.

3.3 Physical Activity

Regular physical activity, either by implementing exercise into daily life activities or by participating in sports, can be considered as an evolutionary based “wonder pill” for humans that has an overall beneficial impact on health throughout life [155, 156]. Fundamental scientific evidence demonstrates that regular physical activity can functionally strengthen and enhance almost every organic system of the human body (e.g. the cardiovascular, respiratory, musculoskeletal, metabolic, neurological, immune, and hematopoietic systems) [157]. Apparently, there is a genetically dependent [158] individual dose response to physical activity that needs to be considered in order to optimize health-related outcomes [159]. On the other hand, a sedentary life style is a well-known risk factor for various diseases and poorer outcomes [160]. In the context of cancer mortality, physical activity is an epidemiologically well-established preventive factor for frequent types of cancers and with a lesser base of evidence, this is true also for other cancers [161, 162]. Furthermore, it is an effective supportive therapeutic strategy during all different phases (pre, during, and post) of cancer therapy even with the potential to lower the risk of cancer reoccurrence [163, 164]. For example, a very recently performed meta-analysis of 26 individual studies found a 37% risk reduction of cancer-specific mortality in the most active patients suffering from either colorectal, breast, or prostate cancer when compared to the least active patients [165]. Considering the previously mentioned aspects of physical activity, it is obvious that the amount of exertion and exercise is a very crucial, if not one of the most important behavioral factors when investigating cancer incidence, prevalence and mortality among populations living at different altitudes. However, in line with a recent review statement by Burtscher *et al.* [48] there is currently a lack of available epidemiological data, which would allow a statistical factorial analysis to evaluate the impact of the level of physical activity, geographical location, and corresponding altitude on cancer-specific mortality. Of note, he recently reported a linear decline of age-standardized mortality rates for male colorectal cancer and female breast cancer with increasing altitude when analyzing data from the Austrian Mortality [166]. The

potentially higher exercise capacity of people living at altitude was proposed, among other factors, to be responsible for the observed lower cancer mortality rate [166]. Driven by the rising socio-economic burden of modern personalized drug-based cancer therapy [167] recent translational efforts across the spectrum from basic to clinical research are beginning to unravel the complex and multifactorial molecular mechanisms responsible for the overall beneficial effects of physical activity on cancer mortality [164]. Obviously, the primary target organs of exercise are the skeletal muscles and heart. Exercise-induced muscle tissue hypoxia may be a key stimulus, predominantly mediated via hypoxia inducible factors (HIFs) signaling cascades [168, 169]. This implicates a complex molecular crosstalk between the two stimuli of physical activity and altitude exposure, which may lead to an altered cellular response in people physically active at altitude. In fact, endurance athletes use various forms of altitude-training (e.g. live high, train high: LHTH) to improve exercise performance by combining both stimuli [170]. In contrast to popular belief, the skeletal musculature is not solely a motor organ but also a highly active endocrine organ that communicates mainly through the vascular circuitry with other body tissues via release of extracellular vesicles and secretory cytokines, known as myokines [171]. A recent preclinical study performed in five different tumor-bearing mouse models found that voluntary exercise on a running-wheel could reduce tumor incidence and growth by more than 60%, especially when applied four weeks before tumor implantation and then continued for the two weeks after tumor implantation [172]. In the same study the authors further reported that these beneficial effects were mainly driven by an epinephrine-dependent mobilization of natural killer cells into the blood stream and interleukin-6 (IL-6)-dependent targeting of these cells to tumors. Interestingly, IL-6 is also a myokine [173], which is known to be secreted from the skeletal muscle upon contraction and probably also hypoxia [174-177]. Another important cellular aspect of physical activity is the fact that exercise can increase the number of adult stem cells (ASCs) [178]. Considering the previously mentioned DeGregori hypothesis of cellular fitness and cancer evolution [47], an exercise-induced increase of healthy ASCs may not only boost tissue regeneration and rejuvenation, but also the anti-oncogenic potential of the corresponding tissue.

To briefly summarize, there is compelling evidence from both animal and human studies that physical activity reduces cancer-specific mortality. However, due to the current lack of

available epidemiological data, no conclusions can be drawn about the impact of physical activity on altered cancer-specific mortality of populations living at different altitudes. Furthermore, the suggested molecular crosstalk between the two stimuli of exercise and altitude exposure further complicates the search for a mechanistic understanding.

4 Conclusion

In many cases, cancer mortality declines with increasing altitude, but the mechanisms that drive tumor suppression are unknown and, due to the complexity of HA adaption processes, difficult to identify. Exposure to HA causes multifactorial cellular, physiological and metabolic alterations that potentially have an impact on tumor mortality. HA and hypoxia induced mechanisms aim at coping with hypoxia and best adapting to HA. Many of these pathways are interconnected in a complex manner. Based on DeGregori's hypothesis of adaptive oncogenesis [47], we suggest that adaptation may increase fitness of healthy cells to better compete with cancer cells within the same niche (Fig.1). In a simplified scheme, one might compare these cells with exercising humans where stress-stimuli increase the cell's fitness level. As a consequence, it is possible that, instead of individual pathways, the sum of all (or some) adaptive processes results in reduced cancer mortality, which may complicate mechanistically studies. However, some feasible studies could shed some more light on this "black box". The poor prognostic factor anemia of cancer with incidence, pathology, severity and response to treatment could usefully contribute to understanding cancer mortality at HA. The recording of physical activity would be desirable in order to compare the activity pattern of "highlander" and "lowlander" cancer patients in relation to mortality. Finally, the efficacy of drugs including their pharmacodynamics and pharmacokinetics needs to be addressed at HA. Drug metabolism is clearly altered in humans exposed to hypoxia and HA and it is fundamental to understand how patients respond to any kind of drug – including cancer therapeutics – in order to guarantee safety and efficacy.

5 Acknowledgements

The authors kindly thank Jeanne Peter for excellent artwork. We acknowledge the financial support of the Swiss National Science Foundation, Marie-Louise von Muralt Foundation, Krebsliga Switzerland and the Zurich Center for Integrative Human Physiology (ZIHP). We apologize to all those authors whose original or review articles could not be cited due to space limitation.

6 References

- [1] Powell FL, Fu Z. HIF-1 and ventilatory acclimatization to chronic hypoxia. *Respir Physiol Neurobiol.* 2008;164:282-7.
- [2] Goldfarb-Rumyantzev AS, Alper SL. Short-term responses of the kidney to high altitude in mountain climbers. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association.* 2014;29:497-506.
- [3] Swenson ER, Olsen NV. Renal Function and Fluid Homeostasis . *High Altitude: Human Adaptation to Hypoxia.* 2014;EDS. Swenson E.R. and Bartsch P.:217-36.
- [4] Haase VH. Regulation of erythropoiesis by hypoxia-inducible factors. *Blood reviews.* 2013;27:41-53.
- [5] Netzer N, Strohl K, Faulhaber M, Gatterer H, Bartscher M. Hypoxia-related altitude illnesses. *Journal of travel medicine.* 2013;20:247-55.
- [6] Smith TG, Robbins PA, Ratcliffe PJ. The human side of hypoxia-inducible factor. *British journal of haematology.* 2008;141:325-34.
- [7] Luks AM, Swenson ER, Bartsch P. Acute high-altitude sickness. *European respiratory review : an official journal of the European Respiratory Society.* 2017;26.
- [8] Semenza GL, Wang GL. A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol Cell Biol.* 1992;12:5447-54.
- [9] Flamme I, Frohlich T, von Reutern M, Kappel A, Damert A, Risau W. HRF, a putative basic helix-loop-helix-PAS-domain transcription factor is closely related to hypoxia-inducible factor-1 alpha and developmentally expressed in blood vessels. *Mech Dev.* 1997;63:51-60.
- [10] Tian H, McKnight SL, Russell DW. Endothelial PAS domain protein 1 (EPAS1), a transcription factor selectively expressed in endothelial cells. *Genes Dev.* 1997;11:72-82.
- [11] Kaelin WG, Jr., Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell.* 2008;30:393-402.
- [12] Hopfl G, Ogunshola O, Gassmann M. HIFs and tumors--causes and consequences. *Am J Physiol Regul Integr Comp Physiol.* 2004;286:R608-23.
- [13] Fandrey J, Gassmann M. Oxygen sensing and the activation of the hypoxia inducible factor 1 (HIF-1)--invited article. *Adv Exp Med Biol.* 2009;648:197-206.
- [14] Kapitsinou PP, Liu Q, Unger TL, Rha J, Davidoff O, Keith B, et al. Hepatic HIF-2 regulates erythropoietic responses to hypoxia in renal anemia. *Blood.* 2010;116:3039-48.
- [15] Hayes DP. Cancer protection related to solar ultraviolet radiation, altitude and vitamin D. *Med Hypotheses.* 2010;75:378-82.
- [16] Brown JM. Tumor hypoxia in cancer therapy. *Methods Enzymol.* 2007;435:297-321.

- [17] Vaupel P, Mayer A. Hypoxia in cancer: significance and impact on clinical outcome. *Cancer Metastasis Rev.* 2007;26:225-39.
- [18] Zhao J, Du F, Shen G, Zheng F, Xu B. The role of hypoxia-inducible factor-2 in digestive system cancers. *Cell Death Dis.* 2015;6.
- [19] Semenza GL. Defining the role of hypoxia-inducible factor 1 in cancer biology and therapeutics. *Oncogene.* 2010;29:625-34.
- [20] Liu ZJ, Semenza GL, Zhang HF. Hypoxia-inducible factor 1 and breast cancer metastasis. *J Zhejiang Univ-Sc B.* 2015;16:32-43.
- [21] Shen C, Beroukhim R, Schumacher SE, Zhou J, Chang M, Signoretti S, et al. Genetic and functional studies implicate HIF1alpha as a 14q kidney cancer suppressor gene. *Cancer discovery.* 2011;1:222-35.
- [22] Lanikova L, Reading NS, Hu H, Tashi T, Burjanivova T, Shestakova A, et al. Evolutionary selected Tibetan variants of HIF pathway and risk of lung cancer. *Oncotarget.* 2016.
- [23] Li K, Zhang Y, Dan Z, Wang Y, Ren ZC. Association of the hypoxia inducible factor-1alpha gene polymorphisms with gastric cancer in Tibetans. *Biochemical genetics.* 2009;47:625-34.
- [24] Simonson TS, Yang Y, Huff CD, Yun H, Qin G, Witherspoon DJ, et al. Genetic evidence for high-altitude adaptation in Tibet. *Science.* 2010;329:72-5.
- [25] Lorenzo FR, Huff C, Myllymaki M, Olenchock B, Swierczek S, Tashi T, et al. A genetic mechanism for Tibetan high-altitude adaptation. *Nature genetics.* 2014;46:951-6.
- [26] Burton AC. Cancer and altitude. Does intracellular pH regulate cell division? *Eur J Cancer.* 1975;11:365-71.
- [27] Eckhoff ND, Shultis JK, Clack RW, Ramer ER. Correlation of leukemia mortality rates with altitude in the United States. *Health Phys.* 1974;27:377-80.
- [28] Mason TJ, Miller RW. Cosmic radiation at high altitudes and U.S. cancer mortality, 1950-1969. *Radiat Res.* 1974;60:302-6.
- [29] Amsel J, Waterbor JW, Oler J, Rosenwaike I, Marshall K. Relationship of site-specific cancer mortality rates to altitude. *Carcinogenesis.* 1982;3:461-5.
- [30] Hart J. Mean cancer mortality rates in low versus high elevation counties in Texas. *Dose Response.* 2010;8:448-55.
- [31] Hart J. Cancer mortality for a single race in low versus high elevation counties in the u.s. *Dose Response.* 2011;9:348-55.
- [32] Hart J. Cancer mortality in six lowest versus six highest elevation jurisdictions in the u.s. *Dose Response.* 2011;9:50-8.
- [33] Hart J, Hyun S. Cancer mortality, state mean elevations, and other selected predictors. *Dose Response.* 2012;10:58-65.
- [34] Weinberg CR, Brown KG, Hoel DG. Altitude, radiation, and mortality from cancer and heart disease. *Radiat Res.* 1987;112:381-90.

- [35] Van Pelt WR. Epidemiological associations among lung cancer, radon exposure and elevation above sea level - A reassessment of Cohen's county level radon study. *Health Phys.* 2003;85:397-403.
- [36] Youk AO, Buchanich JM, Fryzek J, Cunningham M, Marsh GM. An ecological study of cancer mortality rates in high altitude counties of the United States. *High Alt Med Biol.* 2012;13:98-104.
- [37] Boscoe FP, Schymura MJ. Solar ultraviolet-B exposure and cancer incidence and mortality in the United States, 1993-2002. *BMC Cancer.* 2006;6.
- [38] Simeonov KP, Himmelstein DS. Lung cancer incidence decreases with elevation: evidence for oxygen as an inhaled carcinogen. *Peerj.* 2015;2.
- [39] Mori-Chavez P, Upton AC, Salazar M, Conklin JW. Influence of altitude on late effects of radiation in RF-Un mice: observations on survival time, blood changes, body weight, and incidence of neoplasms. *Cancer Res.* 1970;30:913-28.
- [40] Mori-Chavez P. Spontaneous leukemia at high altitude in C58 mice. *Journal of the National Cancer Institute.* 1958;21:985-97.
- [41] Mori-Chavez P. Development of spontaneous pulmonary tumors at high altitude in strain A mice. *Journal of the National Cancer Institute.* 1962;28:55-73.
- [42] Kulish OP. [Characteristics of the metastatic spread of transplanted tumors under high-altitude conditions]. *Eksperimental'naia onkologiya.* 1987;9:73-5.
- [43] Kulish OP. [Effect high-altitude conditions on the antitumor activity of cyclophosphane and its action on lymphoid tissue cells]. *Eksperimental'naia onkologiya.* 1985;7:60-3.
- [44] Kulish OP, Galkina KA. [Effect of high-altitude hypoxia on the effectiveness of chemotherapy in tumors]. *Biulleten' eksperimental'noi biologii i meditsiny.* 1983;95:33-5.
- [45] Sung HJ, Ma WZ, Starost MF, Lago CU, Lim PK, Sack MN, et al. Ambient Oxygen Promotes Tumorigenesis. *Plos One.* 2011;6.
- [46] Germain P. Cancer cells and adaptive explanations. *Biol Philos.* 2012;27:785.
- [47] DeGregori J. Evolved tumor suppression: why are we so good at not getting cancer? *Cancer Res.* 2011;71:3739-44.
- [48] Burtcher M. Effects of Living at Higher Altitudes on Mortality: A Narrative Review. *Aging Dis.* 2014;5:274-80.
- [49] Arnold M, Leitzmann M, Freisling H, Bray F, Romieu I, Renehan A, et al. Obesity and cancer: An update of the global impact. *Cancer Epidemiol.* 2016;41:8-15.
- [50] Goday A, Barneto I, Garcia-Almeida JM, Blasco A, Lecube A, Grivalos C, et al. Obesity as a risk factor in cancer: A national consensus of the Spanish Society for the Study of Obesity and the Spanish Society of Medical Oncology. *Clin Transl Oncol.* 2015;17:763-71.
- [51] Diaz-Gutierrez J, Martinez-Gonzalez MA, Pons Izquierdo JJ, Gonzalez-Muniesa P, Martinez JA, Bes-Rastrollo M. Living at Higher Altitude and Incidence of Overweight/Obesity: Prospective Analysis of the SUN Cohort. *Plos One.* 2016;11:e0164483.

- [52] Woolcott OO, Gutierrez C, Castillo OA, Elashoff RM, Stefanovski D, Bergman RN. Inverse association between altitude and obesity: A prevalence study among andean and low-altitude adult individuals of Peru. *Obesity*. 2016;24:929-37.
- [53] Woolcott OO, Castillo OA, Gutierrez C, Elashoff RM, Stefanovski D, Bergman RN. Inverse association between diabetes and altitude: a cross-sectional study in the adult population of the United States. *Obesity*. 2014;22:2080-90.
- [54] Vander Heiden MG, DeBerardinis RJ. Understanding the Intersections between Metabolism and Cancer Biology. *Cell*. 2017;168:657-69.
- [55] DeBerardinis RJ, Mancuso A, Daikhin E, Nissim I, Yudkoff M, Wehrli S, et al. Beyond aerobic glycolysis: transformed cells can engage in glutamine metabolism that exceeds the requirement for protein and nucleotide synthesis. *Proc Natl Acad Sci U S A*. 2007;104:19345-50.
- [56] Hensley CT, Faubert B, Yuan Q, Lev-Cohain N, Jin E, Kim J, et al. Metabolic Heterogeneity in Human Lung Tumors. *Cell*. 2016;164:681-94.
- [57] Duan W, Shen X, Lei J, Xu Q, Yu Y, Li R, et al. Hyperglycemia, a neglected factor during cancer progression. *BioMed research international*. 2014;2014:461917.
- [58] Augustin LS, Kendall CW, Jenkins DJ, Willett WC, Astrup A, Barclay AW, et al. Glycemic index, glycemic load and glycemic response: An International Scientific Consensus Summit from the International Carbohydrate Quality Consortium (ICQC). *Nutrition, metabolism, and cardiovascular diseases : NMCD*. 2015;25:795-815.
- [59] Bailey DM, Castell LM, Newsholme EA, Davies B. Continuous and intermittent exposure to the hypoxia of altitude: implications for glutamine metabolism and exercise performance. *Br J Sports Med*. 2000;34:210-2.
- [60] Chen L, Cui H. Targeting Glutamine Induces Apoptosis: A Cancer Therapy Approach. *International journal of molecular sciences*. 2015;16:22830-55.
- [61] Woolcott OO, Ader M, Bergman RN. Glucose homeostasis during short-term and prolonged exposure to high altitudes. *Endocrine reviews*. 2015;36:149-73.
- [62] Gerich JE, Meyer C, Woerle HJ, Stumvoll M. Renal gluconeogenesis: its importance in human glucose homeostasis. *Diabetes care*. 2001;24:382-91.
- [63] Labak CM, Wang PY, Arora R, Guda MR, Asuthkar S, Tsung AJ, et al. Glucose transport: meeting the metabolic demands of cancer, and applications in glioblastoma treatment. *American journal of cancer research*. 2016;6:1599-608.
- [64] Wood IS, Wang B, Lorente-Cebrian S, Trayhurn P. Hypoxia increases expression of selective facilitative glucose transporters (GLUT) and 2-deoxy-D-glucose uptake in human adipocytes. *Biochemical and biophysical research communications*. 2007;361:468-73.
- [65] Zoll J, Ponsot E, Dufour S, Doutreleau S, Ventura-Clapier R, Vogt M, et al. Exercise training in normobaric hypoxia in endurance runners. III. Muscular adjustments of selected gene transcripts. *Journal of applied physiology*. 2006;100:1258-66.

- [66] Huang C, Freter C. Lipid metabolism, apoptosis and cancer therapy. *International journal of molecular sciences*. 2015;16:924-49.
- [67] Gajate C, Mollinedo F. Lipid raft-mediated Fas/CD95 apoptotic signaling in leukemic cells and normal leukocytes and therapeutic implications. *Journal of leukocyte biology*. 2015;98:739-59.
- [68] Mollinedo F, Gajate C. Lipid rafts as major platforms for signaling regulation in cancer. *Advances in biological regulation*. 2015;57:130-46.
- [69] Simons K, Toomre D. Lipid rafts and signal transduction. *Nature reviews Molecular cell biology*. 2000;1:31-9.
- [70] Cui Y, Whiteman MK, Flaws JA, Langenberg P, Tkaczuk KH, Bush TL. Body mass and stage of breast cancer at diagnosis. *Int J Cancer*. 2002;98:279-83.
- [71] Caan BJ, Coates AO, Slaterry ML, Potter JD, Quesenberry CP, Jr., Edwards SM. Body size and the risk of colon cancer in a large case-control study. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*. 1998;22:178-84.
- [72] Chow WH, Gridley G, Fraumeni JF, Jr., Jarvholm B. Obesity, hypertension, and the risk of kidney cancer in men. *The New England journal of medicine*. 2000;343:1305-11.
- [73] Bianchini F, Kaaks R, Vainio H. Overweight, obesity, and cancer risk. *Lancet Oncol*. 2002;3:565-74.
- [74] Lowenfels AB, Maisonneuve P, Boyle P, Zatonski WA. Epidemiology of gallbladder cancer. *Hepato-gastroenterology*. 1999;46:1529-32.
- [75] Berrington de Gonzalez A, Sweetland S, Spencer E. A meta-analysis of obesity and the risk of pancreatic cancer. *Br J Cancer*. 2003;89:519-23.
- [76] Hashmi S, Wang Y, Suman DS, Parhar RS, Collison K, Conca W, et al. Human cancer: is it linked to dysfunctional lipid metabolism? *Biochim Biophys Acta*. 2015;1850:352-64.
- [77] Chen J. Multiple signal pathways in obesity-associated cancer. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2011;12:1063-70.
- [78] Fritz V, Fajas L. Metabolism and proliferation share common regulatory pathways in cancer cells. *Oncogene*. 2010;29:4369-77.
- [79] Sears B, Perry M. The role of fatty acids in insulin resistance. *Lipids in health and disease*. 2015;14:121.
- [80] Kamphorst JJ, Cross JR, Fan J, de Stanchina E, Mathew R, White EP, et al. Hypoxic and Ras-transformed cells support growth by scavenging unsaturated fatty acids from lysophospholipids. *Proc Natl Acad Sci U S A*. 2013;110:8882-7.
- [81] Ackerman D, Simon MC. Hypoxia, lipids, and cancer: surviving the harsh tumor microenvironment. *Trends in cell biology*. 2014;24:472-8.
- [82] Acs Z, Bori Z, Takeda M, Osvath P, Berkes I, Taylor AW, et al. High altitude exposure alters gene expression levels of DNA repair enzymes, and modulates fatty acid metabolism by SIRT4 induction in human skeletal muscle. *Respir Physiol Neurobiol*. 2014;196:33-7.

- [83] Allott EH, Hursting SD. Obesity and cancer: mechanistic insights from transdisciplinary studies. *Endocr-Relat Cancer*. 2015;22:R365-R86.
- [84] Gong Y, Dou LJ, Liang J. Link between obesity and cancer: role of triglyceride/free fatty acid cycling. *Eur Rev Med Pharmacol*. 2014;18:2808-20.
- [85] Ranhotra HS, Sharma R. Moderately high altitude habitation modulates lipid profile and alkaline phosphatase activity in aged Khasis of Meghalaya. *Indian journal of clinical biochemistry : IJCB*. 2010;25:51-6.
- [86] Vats P, Ray K, Majumadar D, Amitabh, Joseph DA, Bayen S, et al. Changes in cardiovascular functions, lipid profile, and body composition at high altitude in two different ethnic groups. *High Alt Med Biol*. 2013;14:45-52.
- [87] Kontush A. HDL-mediated mechanisms of protection in cardiovascular disease. *Cardiovascular research*. 2014;103:341-9.
- [88] Kontush A. HDL particle number and size as predictors of cardiovascular disease. *Frontiers in pharmacology*. 2015;6:218.
- [89] Touvier M, Fassier P, His M, Norat T, Chan DS, Blacher J, et al. Cholesterol and breast cancer risk: a systematic review and meta-analysis of prospective studies. *The British journal of nutrition*. 2015;114:347-57.
- [90] Kotani K, Sekine Y, Ishikawa S, Ikpot IZ, Suzuki K, Remaley AT. High-Density Lipoprotein and Prostate Cancer: An Overview. *J Epidemiol*. 2013;23:313-9.
- [91] Mondul AM, Weinstein SJ, Virtamo J, Albanes D. Serum total and HDL cholesterol and risk of prostate cancer. *Cancer Cause Control*. 2011;22:1545-52.
- [92] Su F, Grijalva V, Navab K, Ganapathy E, Meriwether D, Imaizumi S, et al. HDL Mimetics Inhibit Tumor Development in Both Induced and Spontaneous Mouse Models of Colon Cancer. *Mol Cancer Ther*. 2012;11:1311-9.
- [93] Hentze MW, Muckenthaler MU, Galy B, Camaschella C. Two to tango: regulation of Mammalian iron metabolism. *Cell*. 2010;142:24-38.
- [94] Nemeth E, Tuttle MS, Powelson J, Vaughn MB, Donovan A, Ward DM, et al. Heparin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science*. 2004;306:2090-3.
- [95] Muckenthaler MU, Rivella S, Hentze MW, Galy B. A Red Carpet for Iron Metabolism. *Cell*. 2017;168:344-61.
- [96] Gassmann M, Muckenthaler MU. Adaptation of iron requirement to hypoxic conditions at high altitude. *Journal of applied physiology*. 2015;119:1432-40.
- [97] Peyssonnaud C, Nizet V, Johnson RS. Role of the hypoxia inducible factors HIF in iron metabolism. *Cell cycle*. 2008;7:28-32.
- [98] Kautz L, Jung G, Valore EV, Rivella S, Nemeth E, Ganz T. Identification of erythroferrone as an erythroid regulator of iron metabolism. *Nature genetics*. 2014;46:678-84.
- [99] Torti SV, Torti FM. Iron and cancer: more ore to be mined. *Nat Rev Cancer*. 2013;13:342-55.

- [100] Wu T, Sempos CT, Freudenheim JL, Muti P, Smit E. Serum iron, copper and zinc concentrations and risk of cancer mortality in US adults. *Annals of epidemiology*. 2004;14:195-201.
- [101] Nelson RL. Iron and colorectal cancer risk: human studies. *Nutrition reviews*. 2001;59:140-8.
- [102] Recalcati S, Locati M, Marini A, Santambrogio P, Zaninotto F, De Pizzol M, et al. Differential regulation of iron homeostasis during human macrophage polarized activation. *European journal of immunology*. 2010;40:824-35.
- [103] Daniels TR, Bernabeu E, Rodriguez JA, Patel S, Kozman M, Chiappetta DA, et al. The transferrin receptor and the targeted delivery of therapeutic agents against cancer. *Biochim Biophys Acta*. 2012;1820:291-317.
- [104] Kakhlon O, Gruenbaum Y, Cabantchik ZI. Ferritin expression modulates cell cycle dynamics and cell responsiveness to H-ras-induced growth via expansion of the labile iron pool. *The Biochemical journal*. 2002;363:431-6.
- [105] Kakhlon O, Gruenbaum Y, Cabantchik ZI. Repression of ferritin expression modulates cell responsiveness to H-ras-induced growth. *Biochemical Society transactions*. 2002;30:777-80.
- [106] Jiang XP, Elliott RL, Head JF. Manipulation of iron transporter genes results in the suppression of human and mouse mammary adenocarcinomas. *Anticancer research*. 2010;30:759-65.
- [107] Cook JD, Boy E, Flowers C, Daroca Mdel C. The influence of high-altitude living on body iron. *Blood*. 2005;106:1441-6.
- [108] Govus AD, Garvican-Lewis LA, Abbiss CR, Peeling P, Gore CJ. Pre-Altitude Serum Ferritin Levels and Daily Oral Iron Supplement Dose Mediate Iron Parameter and Hemoglobin Mass Responses to Altitude Exposure. *Plos One*. 2015;10:e0135120.
- [109] Kasprzak Z, Sliwicka E, Hennig K, Pilaczynska-Szczesniak L, Huta-Osiecka A, Nowak A. Vitamin D, Iron Metabolism, and Diet in Alpinists During a 2-Week High-Altitude Climb. *High Alt Med Biol*. 2015;16:230-5.
- [110] Yamasaki T, Terai S, Sakaida I. Deferoxamine for advanced hepatocellular carcinoma. *The New England journal of medicine*. 2011;365:576-8.
- [111] Whitnall M, Howard J, Ponka P, Richardson DR. A class of iron chelators with a wide spectrum of potent antitumor activity that overcomes resistance to chemotherapeutics. *Proc Natl Acad Sci U S A*. 2006;103:14901-6.
- [112] Crepin R, Goenaga AL, Jullienne B, Bougherara H, Legay C, Benihoud K, et al. Development of human single-chain antibodies to the transferrin receptor that effectively antagonize the growth of leukemias and lymphomas. *Cancer Res*. 2010;70:5497-506.
- [113] Miller JL. Iron deficiency anemia: a common and curable disease. *Cold Spring Harbor perspectives in medicine*. 2013;3.
- [114] Gilreath JA, Stenehjem DD, Rodgers GM. Diagnosis and treatment of cancer-related anemia. *American journal of hematology*. 2014;89:203-12.

- [115] Krantz SB. Pathogenesis and treatment of the anemia of chronic disease. *Am J Med Sci*. 1994;307:353-9.
- [116] Steinberg D. Anemia and cancer. *CA Cancer J Clin*. 1989;39:296-304.
- [117] Caro JJ, Salas M, Ward A, Goss G. Anemia as an independent prognostic factor for survival in patients with cancer: a systemic, quantitative review. *Cancer*. 2001;91:2214-21.
- [118] Knight K, Wade S, Balducci L. Prevalence and outcomes of anemia in cancer: a systematic review of the literature. *The American journal of medicine*. 2004;116 Suppl 7A:11S-26S.
- [119] Crusz SM, Balkwill FR. Inflammation and cancer: advances and new agents. *Nature reviews Clinical oncology*. 2015;12:584-96.
- [120] Govus AD, Peeling P, Abbiss CR, Lawler NG, Swinkels DW, Laarakkers CM, et al. Live high, train low - influence on resting and post-exercise hepcidin levels. *Scandinavian journal of medicine & science in sports*. 2016.
- [121] Altamura S, Bartsch P, Dehnert C, Maggiorini M, Weiss G, Theurl I, et al. Increased hepcidin levels in high-altitude pulmonary edema. *Journal of applied physiology*. 2015;118:292-8.
- [122] Piperno A, Galimberti S, Mariani R, Pelucchi S, Ravasi G, Lombardi C, et al. Modulation of hepcidin production during hypoxia-induced erythropoiesis in humans in vivo: data from the HIGHCARE project. *Blood*. 2011;117:2953-9.
- [123] Talbot NP, Lakhal S, Smith TG, Privat C, Nickol AH, Rivera-Ch M, et al. Regulation of hepcidin expression at high altitude. *Blood*. 2012;119:857-60.
- [124] Shapiro BB, Streja E, Rhee CM, Molnar MZ, Kheifets L, Kovesdy CP, et al. Revisiting the association between altitude and mortality in dialysis patients. *Hemodialysis international International Symposium on Home Hemodialysis*. 2014;18:374-83.
- [125] Ghahramani N, Ahmed F, Al-Laham A, Lengerich EJ. The epidemiological association of altitude with chronic kidney disease: Evidence of protective effect. *Nephrology*. 2011;16:219-24.
- [126] Brookhart MA, Schneeweiss S, Avorn J, Bradbury BD, Rothman KJ, Fischer M, et al. The effect of altitude on dosing and response to erythropoietin in ESRD. *Journal of the American Society of Nephrology : JASN*. 2008;19:1389-95.
- [127] Sibbel S, Maroni BJ, Brunelli SM. The effect of altitude on erythropoiesis-stimulating agent dose, hemoglobin level, and mortality in hemodialysis patients. *Journal of nephrology*. 2016.
- [128] Henke M, Laszig R, Rube C, Schafer U, Haase KD, Schilcher B, et al. Erythropoietin to treat head and neck cancer patients with anaemia undergoing radiotherapy: randomised, double-blind, placebo-controlled trial. *Lancet*. 2003;362:1255-60.
- [129] Leyland-Jones B. Breast cancer trial with erythropoietin terminated unexpectedly. *Lancet Oncol*. 2003;4:459-60.

- [130] Leyland-Jones B, Semiglazov V, Pawlicki M, Pienkowski T, Tjulandin S, Manikhas G, et al. Maintaining normal hemoglobin levels with epoetin alfa in mainly nonanemic patients with metastatic breast cancer receiving first-line chemotherapy: a survival study. *J Clin Oncol*. 2005;23:5960-72.
- [131] Overgaard J, Hoff C, Hansen HS, Specht L, Overgaard M, Grau C, et al. Randomized study of the importance of Novel Erythropoiesis Stimulating Protein (Aranesp) for the effect of radiotherapy in patients with primary squamous cell carcinoma of the head and neck (HNSCC) — the Danish Head and Neck Group DAHANCA 10 randomized trial. *Eur J Cancer Suppl* 5. 2007;7.
- [132] Smith RE, Jr., Aapro MS, Ludwig H, Pinter T, Smakal M, Ciuleanu TE, et al. Darbepoetin alpha for the treatment of anemia in patients with active cancer not receiving chemotherapy or radiotherapy: results of a phase III, multicenter, randomized, double-blind, placebo-controlled study. *J Clin Oncol*. 2008;26:1040-50.
- [133] Temkin SM, Hellmann M, Serur E, Lee YC, Abulafia O. Erythropoietin administration during primary treatment for locally advanced cervical carcinoma is associated with poor response to radiation. *Int J Gynecol Cancer*. 2006;16:1855-61.
- [134] Thomas G, Ali S, Hoebbers FJ, Darcy KM, Rodgers WH, Patel M, et al. Phase III trial to evaluate the efficacy of maintaining hemoglobin levels above 12.0 g/dL with erythropoietin vs above 10.0 g/dL without erythropoietin in anemic patients receiving concurrent radiation and cisplatin for cervical cancer. *Gynecol Oncol*. 2008;108:317-25.
- [135] Wright JR, Ung YC, Julian JA, Pritchard KI, Whelan TJ, Smith C, et al. Randomized, double-blind, placebo-controlled trial of erythropoietin in non-small-cell lung cancer with disease-related anemia. *J Clin Oncol*. 2007;25:1027-32.
- [136] Jelkmann W, Bohlius J, Hallek M, Sytkowski AJ. The erythropoietin receptor in normal and cancer tissues. *Crit Rev Oncol Hematol*. 2008;67:39-61.
- [137] Acs G, Acs P, Beckwith SM, Pitts RL, Clements E, Wong K, et al. Erythropoietin and erythropoietin receptor expression in human cancer. *Cancer Res*. 2001;61:3561-5.
- [138] Acs G, Zhang PJ, McGrath CM, Acs P, McBroom J, Mohyeldin A, et al. Hypoxia-inducible erythropoietin signaling in squamous dysplasia and squamous cell carcinoma of the uterine cervix and its potential role in cervical carcinogenesis and tumor progression. *Am J Pathol*. 2003;162:1789-806.
- [139] Szenajch J, Wcislo G, Jeong JY, Szczylik C, Feldman L. The role of erythropoietin and its receptor in growth, survival and therapeutic response of human tumor cells From clinic to bench - a critical review. *Biochim Biophys Acta*. 2010;1806:82-95.
- [140] Hardee ME, Cao Y, Fu P, Jiang X, Zhao Y, Rabbani ZN, et al. Erythropoietin blockade inhibits the induction of tumor angiogenesis and progression. *Plos One*. 2007;2:e549.
- [141] Hardee ME, Kirkpatrick JP, Shan S, Snyder SA, Vujaskovic Z, Rabbani ZN, et al. Human recombinant erythropoietin (rEpo) has no effect on tumour growth or angiogenesis. *Br J Cancer*. 2005;93:1350-5.

- [142] Katz O, Barzilay E, Skaat A, Herman A, Mittelman M, Neumann D. Erythropoietin induced tumour mass reduction in murine lymphoproliferative models. *Acta Haematol.* 2005;114:177-9.
- [143] Mittelman M, Neumann D, Peled A, Kanter P, Haran-Ghera N. Erythropoietin induces tumor regression and antitumor immune responses in murine myeloma models. *Proc Natl Acad Sci U S A.* 2001;98:5181-6.
- [144] Yasuda Y, Fujita Y, Matsuo T, Koinuma S, Hara S, Tazaki A, et al. Erythropoietin regulates tumour growth of human malignancies. *Carcinogenesis.* 2003;24:1021-9.
- [145] Administration UFaD. Continuing reassessment of the risks of erythropoiesis-stimulating agents (ESAs) administered for the treatment of anemia associated with cancer chemotherapy. FDA briefing document, Oncologic Drugs Advisory Committee 2007. 2007:<http://www.fda.gov/ohrms/dockets/ac/07/briefing/2007-4301b2-02-FDA.pdf>.
- [146] Administration UFaD. Information for Healthcare Professionals: Erythropoiesis Stimulating Agents (ESA) [Aranesp (darbepoetin), Epogen (epoetin alfa), and Procrit (epoetin alfa)]. 2007: <https://http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientandProviders/ucm126481.htm>.
- [147] Bailey DM. On the Significance of Altered Drug Pharmacokinetics-Pharmacodynamics at High Altitude. *High Alt Med Biol.* 2017.
- [148] Cohen JE, Small C. Hypsographic demography: the distribution of human population by altitude. *Proc Natl Acad Sci U S A.* 1998;95:14009-14.
- [149] Conklin KA. Chemotherapy-associated oxidative stress: impact on chemotherapeutic effectiveness. *Integrative cancer therapies.* 2004;3:294-300.
- [150] Okon IS, Zou MH. Mitochondrial ROS and cancer drug resistance: Implications for therapy. *Pharmacological research.* 2015;100:170-4.
- [151] Semenza GL. Oxygen-dependent regulation of mitochondrial respiration by hypoxia-inducible factor 1. *The Biochemical journal.* 2007;405:1-9.
- [152] Jacobs RA, Siebenmann C, Hug M, Toigo M, Meinild AK, Lundby C. Twenty-eight days at 3454-m altitude diminishes respiratory capacity but enhances efficiency in human skeletal muscle mitochondria. *Faseb J.* 2012;26:5192-200.
- [153] Murray AJ, Horscroft JA. Mitochondrial function at extreme high altitude. *J Physiol.* 2016;594:1137-49.
- [154] du Souich P, Fradette C. The effect and clinical consequences of hypoxia on cytochrome P450, membrane carrier proteins activity and expression. *Expert opinion on drug metabolism & toxicology.* 2011;7:1083-100.
- [155] Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation.* 2007;116:1081-93.
- [156] Lee IM, Buchner DM. The importance of walking to public health. *Med Sci Sports Exerc.* 2008;40:S512-8.

- [157] Garatachea N, Pareja-Galeano H, Sanchis-Gomar F, Santos-Lozano A, Fiuza-Luces C, Moran M, et al. Exercise attenuates the major hallmarks of aging. *Rejuvenation research*. 2015;18:57-89.
- [158] Zadro JR, Shirley D, Andrade TB, Scurrah KJ, Bauman A, Ferreira PH. The Beneficial Effects of Physical Activity: Is It Down to Your Genes? A Systematic Review and Meta-Analysis of Twin and Family Studies. *Sports Med Open*. 2017;3:4.
- [159] Wasfy MM, Baggish AL. Exercise Dose in Clinical Practice. *Circulation*. 2016;133:2297-313.
- [160] Tremblay MS, Colley RC, Saunders TJ, Healy GN, Owen N. Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab*. 2010;35:725-40.
- [161] Friedenreich CM, Neilson HK, Lynch BM. State of the epidemiological evidence on physical activity and cancer prevention. *Eur J Cancer*. 2010;46:2593-604.
- [162] Lee IM. Physical activity and cancer prevention--data from epidemiologic studies. *Med Sci Sports Exerc*. 2003;35:1823-7.
- [163] Buffart LM, Kalter J, Sweegers MG, Courneya KS, Newton RU, Aaronson NK, et al. Effects and moderators of exercise on quality of life and physical function in patients with cancer: An individual patient data meta-analysis of 34 RCTs. *Cancer Treat Rev*. 2017;52:91-104.
- [164] Rogers CJ, Colbert LH, Greiner JW, Perkins SN, Hursting SD. Physical activity and cancer prevention : pathways and targets for intervention. *Sports Med*. 2008;38:271-96.
- [165] Friedenreich CM, Neilson HK, Farris MS, Courneya KS. Physical Activity and Cancer Outcomes: A Precision Medicine Approach. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2016;22:4766-75.
- [166] Bartscher M. Lower mortality rates in those living at moderate altitude. *Aging*. 2016;8:2603-4.
- [167] Luengo-Fernandez R, Leal J, Gray A, Sullivan R. Economic burden of cancer across the European Union: a population-based cost analysis. *Lancet Oncol*. 2013;14:1165-74.
- [168] Ameln H, Gustafsson T, Sundberg CJ, Okamoto K, Jansson E, Poellinger L, et al. Physiological activation of hypoxia inducible factor-1 in human skeletal muscle. *Faseb J*. 2005;19:1009-11.
- [169] Lindholm ME, Rundqvist H. Skeletal muscle hypoxia-inducible factor-1 and exercise. *Exp Physiol*. 2016;101:28-32.
- [170] Khodaei M, Grothe HL, Seyfert JH, VanBaak K. Athletes at High Altitude. *Sports health*. 2016;8:126-32.
- [171] Schnyder S, Handschin C. Skeletal muscle as an endocrine organ: PGC-1 α , myokines and exercise. *Bone*. 2015;80:115-25.
- [172] Pedersen L, Idorn M, Olofsson GH, Lauenborg B, Nookaew I, Hansen RH, et al. Voluntary Running Suppresses Tumor Growth through Epinephrine- and IL-6-Dependent NK Cell Mobilization and Redistribution. *Cell Metab*. 2016;23:554-62.

- [173] Pal M, Febbraio MA, Whitham M. From cytokine to myokine: the emerging role of interleukin-6 in metabolic regulation. *Immunol Cell Biol.* 2014;92:331-9.
- [174] Mazzeo C, Swenson ER. The Immune System. *High Altitude: Human Adaptation to Hypoxia.* 2014;EDS. Swenson E.R. and Bartsch P.:271-84.
- [175] Pedersen BK. Muscular interleukin-6 and its role as an energy sensor. *Med Sci Sports Exerc.* 2012;44:392-6.
- [176] Pedersen BK, Fischer CP. Beneficial health effects of exercise--the role of IL-6 as a myokine. *Trends Pharmacol Sci.* 2007;28:152-6.
- [177] Pedersen BK, Steensberg A. Exercise and hypoxia: effects on leukocytes and interleukin-6-shared mechanisms? *Med Sci Sports Exerc.* 2002;34:2004-13.
- [178] Macaluso F, Myburgh KH. Current evidence that exercise can increase the number of adult stem cells. *Journal of muscle research and cell motility.* 2012;33:187-98.




Figure legend:

Figure 1: How high altitude exposure might influence cancer cells.

Assuming that tumorigenesis is an evolutionary selection process, it requires cancer cells (violet) to compete with healthy cells (orange) within the same, resource-restricted niche. If cancer cells are able to successfully populate this niche mainly depends on the quality of the microenvironment (e.g. nutrition availability) and how effective healthy cells are able to compete against cancer cells. In other words, healthy cells with a high cellular fitness as well as an increased demand for nutrition are better competitors and thus impede more successfully the tumor development. Based on this assumption, we suggest a hypothesis for reduced cancer mortality at high altitude: At low altitude (normoxia) expanding tumors increase expression of HIFs that partially contribute to an elevated uptake of lipids, glucose and iron that are essential for tumor progression. Healthy cells in contrast display no increase in HIF-dependent gene expression at low altitude. With increasing altitude however, healthy cells experience a hypoxia-driven, partially HIF-dependent shift in cellular metabolism (similar to tumor cells at low altitude) that may lead to increased nutrition demand. With the activation of nutrition uptake and utilization pathways healthy cells are more competitive, and thus fitter, against cancer cells. Moreover, high altitude exposure leads to a reduction of glucose as well as lipids and potentially iron plasma levels. This suggests that in addition to higher uptake rates of healthy cells the general availability of resources is declining with increasing altitude resulting in unfavorably environmental conditions for tumor growth and thus in a reduction of cancer cell fitness. On top of that, physical activity (e.g. exercise), that is already tumor-suppressive at low altitude may be boosted in combination with HA exposure.